

Chloroquine

Help

Main

New Search

Search Results

DrugPoints® System

CHLOROQUINE PHOSPHATE

- **Common Tradenames (See Complete Tradename Listing)**

- ARALEN

- **Class**

- antimalarial

- **Dosage, Adult (usual)**

- Amebiasis: 1000 mg chloroquine phosphate (600 mg base) ORALLY daily for 2 days, then 500 mg chloroquine phosphate (300 mg base) ORALLY daily for 2-3 weeks
- Malaria treatment, uncomplicated acute attacks: 1000 mg chloroquine phosphate (600 mg base) ORALLY, then 500 mg chloroquine phosphate (300 mg base) ORALLY after 6-8 hours, then 500 mg chloroquine phosphate (300 mg base) ORALLY once daily for 2 consecutive days; total dose 2500 mg chloroquine phosphate (1500 mg base)
- Malaria treatment, severe: 10 mg base/kg IV infused over 8 hours, then 15 mg base/kg infused over 24hrs; oral therapy should be substituted as soon as patient can tolerate oral medication
- Malaria suppression: 500 mg chloroquine phosphate (300 mg base) ORALLY once weekly (on same day of the week); therapy should begin 2wks before and continue 8wks after last exposure to endemic area; if therapy cannot begin 2 weeks before exposure, an initial loading dose of 1000 mg chloroquine phosphate (600 mg base) ORALLY should be administered
- Malaria treatment, severe: initially, 160-200 mg base IM and repeated in 6 hours if necessary, a total dose of 800 mg in the first 24 hours should not be exceeded; oral therapy should be substituted as soon as the patient can tolerate oral medication for a total course of 1.5 grams in 3 days
- Amebiasis: 160-200 mg base IM daily for 10-12 days

- **Dosage, Pediatric, (usual)**

- Malaria treatment: (infants and children) first dose, 10 mg base/kg (maximum of 600 mg base) ORALLY; second dose 6 hours after first dose, 5 mg base/kg (maximum of 300 mg base); third dose 18 hours after second dose, 5 mg base/kg; fourth dose 24 hours after third dose, 5 mg base/kg
- Malaria suppression: (children) 5 mg base/kg (maximum of 300 mg base) ORALLY once weekly on the same day of the week beginning 2 weeks before and continuing for 8 weeks after last exposure in endemic area
- Malaria treatment: extreme caution is advised, 5 mg base/kg mg base IM and repeated in 6 hours if necessary, a total dose of 10 mg base/kg in any 24 hour period should not be exceeded, maximum single dose is 5 mg base/kg; oral therapy should be substituted as soon as the patient can tolerate oral medication
- **Dose Adjustments:**
 - renal impairment: CrCl less than 10 mL/min, 50% of dose; if prolonged therapy is needed, give 50 to 100 mg base/day
 - liver disease: serum drug monitoring may be necessary; 30-50% of dose is modified by the liver
- **Administration**
 - 300 mg base=500 mg Chloroquine Phosphate
 - IV use of chloroquine phosphate is not recommended in children
- **Monitoring**
 - CBC periodically
 - liver and renal function
 - periodic ophthalmologic exams
- **How Supplied**
 - 50 MG/ML SOLUTION FOR INJECTION
 - 250 MG, 500 MG TABLET
- **Indications**
 - **FDA labeled indications**
 - Amebiasis, extraintestinal
 - Malaria
- **Contraindications**
 - hypersensitivity to 4-aminoquinoline compounds
 - retinal/visual field changes
- **Precautions**
 - liver disease, alcoholism, or concurrent administration with known hepatotoxic drugs

- blood dyscrasias
- glucose 6-phosphate dehydrogenase deficiency
- renal impairment
- psoriasis or porphyria
- **Adverse Effects**
 - **COMMON**
 - amnesia
 - ECG changes
 - methemoglobinemia (rare)
 - muscle weakness
 - nausea, vomiting, anorexia, diarrhea, abdominal cramps
 - pruritus
 - QT interval prolongation
 - retinopathy
- **Drug Interactions**
 - aurothioglucose
 - cimetidine
 - droperidol
 - erythromycin
 - foscarnet
 - halofantrine
 - isradipine
 - kaolin
 - levomethadyl
 - magaldrate
 - magnesium carbonate
 - magnesium hydroxide
 - magnesium oxide
 - magnesium trisilicate
 - mefloquine
 - rabies vaccine
 - ziprasidone
- **Pregnancy Category**
 - C
- **Breast Feeding**
 - safe

Healthcare Series Vol. 119 expires 3/2004

- Content for use only by healthcare professionals in conjunction with clinical data. See complete Warranties and disclaimers.

Help

Main

New Search

Search Results

Sinemet

Help

Main

New Search

MICROMEDEX(R) Healthcare Series Integrated Index

Terms Matched{SINEMET; }

• **Summary Documents**

- [Drug Summary Information \[CARBIDOPA/LEVODOPA - Drug Summary Information\]](#)

• **Drug Information**

- [Ingredients from DRUGDEX Tradename Products \[4 Related Occurrences\]](#)
- [PHYSICIANS' DESK REFERENCE \[SINEMET CR TABLETS - Complete Monograph\]](#)
- [Ingredients from MARTINDALE Tradename Products \[2 Related Occurrences\]](#)
- [INDEX NOMINUM \[2 Related Occurrences\]](#)
- [USP DI\(R\) Drug Information for the Health Care Professional \[CARBIDOPA AND LEVODOPA \(SYSTEMIC\)\]](#)
- [USP DI\(R\) Advice for the Patient\(R\) \[LEVODOPA \(SYSTEMIC\)\]](#)
- [▶ List Of DRUGDEX® Tradename Products](#)
- [▶ List Of MARTINDALE Tradename Products](#)

• **Toxicology Information**

- [▶ List Of POISINDEX® Products](#)
- [▶ IDENTIDEX® System Tablet & Capsule Identification](#)

• **Complementary and Alternative Medicine**

- [AltMed-REAX\(TM\) for the Professional \[5-HTP AND CARBIDOPA - Complete Monograph\]](#)

• **Reproductive Risk**

- [TERIS \[LEVODOPA\]](#)

You may modify your search: Sinemet

Search

☐ Find exact term only[Return to TOP of this page](#)

© 1974 - 2004 Thomson MICROMEDEX. All rights reserved. MICROMEDEX(R) Healthcare Series Vol. 119 expires 3/2004. USPDI / Advice for the Patient are registered USP trademarks used herein under license.

Help

Main

New Search

Levodopa

Help

Main

New Search

Search Results

DrugPoints® System

CARBIDOPA/LEVODOPA**• Common Tradenames (See Complete Tradename Listing)**

- ATAMET
- SINEMET 10-100
- SINEMET 25-100
- SINEMET 25-250
- SINEMET CR

• Class

- antiparkinsonian
- antiparkinsonian, dopaminergic

• Dosage, Adult (usual)

- Parkinson's disease: 25/100 TAB; initial, 1 TAB ORALLY 3 times a day; increase by 1 TAB daily or every other day to 8 TABS daily
- Parkinson's disease: 10/100 TAB; initial, 1 TAB ORALLY 3 or 4 times daily; increase by 1 TAB daily or every other day to 8 TABS daily
- Parkinson's disease: 50/200 (sustained release) TAB, 1 TAB ORALLY twice daily at an interval of at least 6 hr
- Parkinson's disease: (sustained release) titration, doses and dosing intervals may be increased or decreased depending upon therapeutic response; most patients adequately treated with doses that provide 400-1600 mg of levodopa per day, administered as divided doses at intervals of 4-8 hr during the waking day; dosage adjustment interval at least 3 days
- Parkinson's disease: maintenance, individualize; minimum of 70-100 mg carbidopa daily to minimize nausea and vomiting, MAX 200 mg carbidopa daily
- Parkinson's disease: conversion from levodopa monotherapy, levodopa should be discontinued at least 12 hr prior to initiation of treatment with carbidopa/levodopa, which daily dose should be 25% of the previous levodopa dosage

- Restless leg syndrome: 25/100 TAB; 1 TAB once daily at bedtime, may repeat dose if awakening within 2 hr
- Restless leg syndrome: 50/200 sustained release TAB; 1 or 2 TAB 1 hr before bedtime
- **Dosage, Pediatric, (usual)**
 - safety and efficacy not established in pediatric patients
- **Administration**
 - do NOT chew or crush sustained release tablets
 - when doses of the sustained release tablets are given at intervals of less than 4 hr, and/or if divided doses are not equal, it is recommended that the smaller doses be given at the end of the day
- **Monitoring**
 - periodic evaluations of hepatic, hematopoietic, cardiovascular, and renal function during extended therapy
 - monitor closely during dose adjustment period with regard to appearance or worsening of involuntary movements, dyskinesia, or nausea
 - observe carefully for symptoms resembling neuroleptic malignant syndrome if abrupt reduction or discontinuation is required, especially if the patient is receiving neuroleptics
- **How Supplied**
 - 10 MG-100 MG, 25 MG-100 MG, 25 MG-250 MG TAB
 - 25 MG-100 MG, 50 MG-200 MG TER
- **Indications**
 - **FDA labeled indications**
 - Parkinson's disease; idiopathic, post-encephalitic parkinsonism, and symptomatic parkinsonism
 - **Non-FDA labeled indications**
 - Restless leg syndrome
- **Contraindications**
 - hypersensitivity to levodopa/carbidopa products
 - history of melanoma, undiagnosed skin lesions
 - narrow-angle glaucoma
 - nonselective MAO inhibitors concurrently or less than 2 wks prior
- **Precautions**
 - abrupt discontinuation/dose reduction of carbidopa/levodopa (risk of neuroleptic malignant-like syndrome, particularly in patients receiving neuroleptics; gradual tapering of the dose is indicated)
 - asthmatic patients or other severe pulmonary disease (potential

exacerbation due to adverse respiratory side effects)

- CNS adverse effects may occur sooner during carbidopa/levodopa therapy than with levodopa alone)
- dose reduction may be indicated if dyskinesias occur during therapy
- hepatic insufficiency (potential exacerbation)
- history of peptic ulcer disease (risk of gastrointestinal bleeding recurrence)
- patients with endocrine diseases/disorders (potential for adverse effects of levodopa on hypothalamic or pituitary function)
- renal impairment (potential for urinary retention)
- residual atrial, nodal, or ventricular arrhythmias following myocardial infarction (potential exacerbation)
- severe cardiovascular disease (arrhythmia potential)
- chronic wide-angle glaucoma (potential for slight increase in intraocular pressure)
- concomitant use of tricyclic antidepressants, dopamine D2 receptor antagonists
- concurrent antihypertensive therapy (postural hypertension risk)
- diabetes mellitus (potential changes in blood-glucose control)
- history of melanoma
- underlying depression or psychosis (potential exacerbation and enhanced suicidal risk)

- **Adverse Effects**

- **COMMON**

- anorexia, nausea, vomiting

- **SERIOUS**

- cardiac abnormalities, orthostatic hypotension (1%)
 - dyskinesias (frequent)
 - psychotic symptoms


- **Drug Interactions**

- acetophenazine
 - bromperidol
 - bupropion
 - chlorpromazine
 - clorgyline
 - droperidol
 - ferric ammonium citrate
 - fluphenazine
 - haloperidol

- iproniazid
- iron
- isocarboxazid
- isoniazid
- mesoridazine
- methotrimeprazine
- metoclopramide
- moclobemide
- nialamide
- papaverine
- pargyline
- perphenazine
- phenelzine
- phenytoin
- pipotiazine
- procarbazine
- prochlorperazine
- promazine
- propiomazine
- pyridoxine
- risperidone
- selegiline
- spiramycin
- thiethylperazine
- thioridazine
- toloxatone
- tranylcypromine
- trifluoperazine
- triflupromazine
- zotepine
- **Pregnancy Category**
 - C
- **Breast Feeding**
 - unknown
- **Notes**
 - when discontinuing levodopa/carbidopa therapy, gradual tapering of the dose is indicated to prevent the occurrence of a condition resembling neuroleptic malignant syndrome
 - may alter some liver function tests, blood urea nitrogen, and positive

Coombs test

- may cause false positive reaction for urinary ketone bodies when a test tape is used for determination of ketonuria
- may cause false-negative tests with the use of glucose-oxidase methods of testing for glucosuria
- rare reports of falsely diagnosed pheochromocytoma

 © 1974 - 2004 Thomson MICROMEDEX. All rights reserved. MICROMEDEX(R) Healthcare Series Vol. 119 expires 3/2004
- Content for use only by healthcare professionals in conjunction with clinical data. See complete Warranties and disclaimers.

[Help](#)[Main](#)[New Search](#)[Search Results](#)